










AT

**Acyl peptides and pharmaceutical preparations thereof, as well as their manufacture and use.**






**Patent number:** EP0030920  
**Publication date:** 1981-06-24  
**Inventor:** RINK HANS; KAMBER BRUNO DR; SIEBER PETER  
**Applicant:** CIBA GEIGY AG (CH)  
**Classification:**  
- international: C07C103/52; A61K37/02  
- european: C07K14/655A  
**Application number:** EP19800810381 19801208  
**Priority number(s):** CH19790011096 19791214

**Also published as:**

 US4369179 (A1)  
 JP56092850 (A)  
 FI803849 (A)  
 ES8200861 (A)  
 EP0030920 (A3)  
 EP0030920 (B1)  
 PT72191 (B)  
 GR73621 (B)  
 DD154098 (B)

less &lt;&lt;

**Cited documents:**

 DE2659758  
 US4113714  
 US4118380  
 EP0011366  
 US4130554  
more >>

Abstract not available for EP0030920

Abstract of corresponding document: **US4369179**

Acylpeptides derived from somatostatin or from a derivative thereof having analogous action, in which the amino acids sequence is modified by omitting individual amino acids and/or by exchanging them for other amino acids, and in which the epsilon -amino group of the lysine residue in the 9-position, and optionally also the epsilon -amino group of the lysine residue in the 4-position and/or the N-terminal alpha -amino group carries the acyl radical of an optionally substituted alkanecarboxylic acid, and salts and complexes thereof can be used as antidiabetics and/or for combating gastrointestinal bleeding. They are manufactured by conventional methods of peptide chemistry.

---

Data supplied from the **esp@cenet** database - Worldwide